









Management Tools to Increase Dairy Cow Feed Efficiency

V.E. Cabrera and A. Kalantari


University of Wisconsin-Madison Dairy Science











This site is designed to support dairy farming decision-making focusing on model-based scientific research. The ultimate goal is to provide user-friendly computerized decision support tools to help dairy farmers improve their economic performance along with environmental stewardship.



University of Wisconsin

University of Wisconsin - Madison **UW - Cooperative Extension** UW - Dairy Science **Dairy Cattle Reproduction Dairy Cattle Nutrition** Milk Quality **UW Dairy Nutrient Understanding Dairy Markets UW Center for Dairy Profitability**

Latest Projects

Improving Dairy Farm Sustainability Genomic Selection and Herd Management Dairy Reproduction Decision Support Tools Strategies of Pasture Supplementation Improving Dairy Cow Fertility

Contact



Associate Professor Extension Specialist in Dairy Management 279 Animal Sciences 1675 Observatory Dr. Madison, WI 53706 (608) 265-8506 vcabrera@wisc.edu More »

Victor E.Cabrera, Ph.D.









Helpful Link

Repro Money Program
















Tools

A collection of the state-of-the-art and scientific-based dairy farm management decision support tools that are user-friendly, interactive, robust, visually attractive, and self-contained. These tools count with associated documentation and video demonstrations. Technical support on their application is also available upon request.

Feeding

- > FeedVal 2012
- > Grouping Strategies for Feeding Lactating Dairy Cattle
- Nutritional Grouping in Wisconsin and Michigan Dairy Farms
- Ontigen® Evaluator

Heifers

Reproduction

Genomics

Production

+40 Decision Support Tools

Replacement

Health

Financial

Feeding all lactating cows equally

A larger number of cows are overfed

Same ration (TMR) to all cows (groups)

All lactating cows receive same nutrient density diet



Preferred "high" rations

Low producing animals receive more nutrients than required

One diet for all Would never optimize production and efficiency

Considering nutritional grouping

For improved feed efficiency

Opportunity to increase productivity

Cows receive more precise diets

Improved profitability


IOFC gains far exceed additional expenses or losses

Diets closer to requirements

Saves feed costs and increases income over feed costs

Additional benefits


- + environmental
 concerns Wang et al., 2000
- † health conditions



Why farmers do not group more?

Trying to find most important constraints

2-page mailed survey




Results (responses)

- 196 WI farms
- 211 MI farms

Constraints to feeding more ration groups

- 1. Milk drops when cows are moved
- 2. Desire to keep management simple
- 3. Conflicts with grouping for reproduction
- 4. Farm facilities do not allow it
- 5. Not enough labor or personnel to handle it



Strategies for grouping cows

Depend on farm and herd characteristics

Individual cow nutrient requirements

- Energy
- Protein (RUP, RDP, MP)

Number of lactating cows on the herd

States (i.e., current characteristics of the cow)

Farm characteristics

Capacity to handle lactating feeding groups



Adapted from McGilliard et al., 1983; St-Pierre and Thraen, 1999

Milk (and components)

Cow-specific lactation curves

Milk based on


- Herd ME305
- Cow PPA or ME305
- Stochasticity

Components

- Herd
- Stochasticity

Base function

- Woods
- Adjusted Woods



De Vries, 2001

Initial individual cow BW


Cow-specific BW



- 1. Available from farm records, or
- 2. Stochastic distribution

Daily BW and BCS change according to:

- Lactation
- •DIM
- Stochasticity



Days after calving

Criteria for nutritional grouping

Several criteria exist

Days after calving (DIM)

Based on stage of lactation



Fat (protein) corrected milk


Based on level of production measured as F(P)CM

Dairy merit

Function of both F(P)CM and BW

Cluster

Seems to be MOST efficient criterion



McGilliard et al., 1983 St-Pierre and Thraen, 1999

Nutritional grouping


Two main types of groups


Obligated groups

- Fresh (< 22 DIM)
- Dry (~> 220 DCC)
- Daily assigned

Optional groups

- Actual additional groups
- Daily assigned
- Monthly re-grouped





Cow and herd simulation

Monte Carlo approach

Next event scheduling

- Pregnancy
- Abortion
- Dry-off
- Parturition
- Involuntary culling
- Death

Two-step

- •1. Binary outcome of event:
 - Happens or not
 - E.g., uniform distribution
- •2. DIM of the occurrence
 - When it happens
 - E.g., Weibull distribution

Immediate replacement

After a cow leaves the herd

Replicates

 1,000 replicates for each cow within specific herd

Cow simulation

Follows actual COW card

Variable	Unit	Description
Cow ID	#	Cow identification
Parity	#	Lactation
DIM	d	Days in milk, days after calving
DCC	d	Days in pregnancy (DIP)
Fat	%	Fat component on milk
Protein	%	Protein component on milk (%)
PPA*	%	Predicted producing ability
ME 305*	kg/305 d	Mature equivalent milk production
BW	kg	Live body weight

^{*}Either PPA or ME305 used to assess cow's milk class. PPA preferred if available


Studied herds

All data collected at the cow-level


Herd (size)	570	787	727	331	1460
Herd ME 305, kg	16,140	12,884	13,897	13,348	14,188
1 st lactation, %	43	39	39	38	45
Average DIM	187	178	201	208	189
21-d PR, %	18	19	19	17	18
Culling risk, %	32	37	36	35	40
Abortion, %	7	11	11	16	7
BW available	X	X	√	√	X




Energy requirements of cows




Energy provided in diets



Energy and Protein concentrations throughout Lactations (1, 2, ≥ 3)




Provided - Required Energy in diet




592 lactating cows (all lactating - fresh) from 787-cow herd

Provided - Required MP in diet



592 lactating cows (all lactating - fresh) from 787-cow herd

Average gain of grouping




Economic Gain (\$/cow.yr)


Farm and	Scenario -	Difference between Grouping and 1 Group		
Herd Size	Scenario	2 Groups	3 Groups	4 Groups
	base	53.54	65.91	-
331	milk loss ¹	35.75	47.59	_
	1st lactation ²	47.66	58.29	_
	base	54.0	65.79	_
570	milk loss ¹	37.59	44.82	_
	1st lactation ²	43.28	53.45	-
	base	62.72	74.98	-
727	milk loss ¹	49.63	54.75	_
	1 st lactation ²	49.89	59.47	-
	base	73.50	88.41	-
787	milk loss ¹	57.53	67.39	-
	1st lactation ²	61.80	74.64	-
	base	57.57	69.96	74.45
1,460	milk loss ¹	43.56	49.36	50.81
	1 st lactation ²	46.90	57.19	61.45

¹1.82kg x 5 d. ²1st lactation fed as a separate group


Energy captured in milk



Nitrogen captured in milk



Body weight and BCS



1,000 replicates for 787-cow herd

Total are under curves adds to 1




Decision support tool...

http://DairyMGT.info

A simplified online tool

Herd-specific assessments (DairyMGT.info)



Additional costs and benefits

Impacts grouping feeding strategies

Management cost

- Additional labor
- Extra management

Avoid costs

 Additives and supplements savings


Milk depression

Cow social interactions



Grouping Strategies

Farm/herd possibilities and decision-making





Tool demonstration

Grouping Illustration

Economic impact of nutritional grouping

Current Situation				
Lactating	470			
Current Groups	None			
NEL Mcal/lb	0.80			
CP, %	17			

Possible Situation				
Groups	3			
Group Sizes	100, 100, 270			
Milk loss	2.27 kg/d x 4 d			
Added Costs	\$1,000/month			
Saved	None			

Decision Support System Illustration

Cluster grouping criteria

Current Situation						
Group	Group Cows NEL CP IOFC					
	#	Mcal/lb	%	\$/cow.d		
All	470	0.80	17.00	6.9		



Annual value of grouping \$135,000/herd

Possible Situation						
Group	Cows	NEL	CP	IOFC		
	#	Mcal/lb	%	\$/cow.d		
1	100	0.62	13.07	4.7		
2	100	0.65	14.18	7.2		
3	270	0.71	16.05	9.3		
All	470	0.68	15.02	7.9		

Wisconsin herds analysis



Analysis from dairy farm records

30 Wisconsin dairy farms

No grouping vs. 3 groups

Same size groups

Grouping criterion

Cluster



Same prices for all

- \$0.35/kg milk
- •\$0.315/kg CP
- \$0.1174/Mcal NE_L

Projected body weight

- 500 kg primiparous
- 600 kg multiparous

Analysis from dairy farm records

30 Wisconsin dairy farms

	Lactating cows (n=30)	1 Group	3 Groups	Gain
		Income Over Feed Cost \$/cow.yr		
Minimum	<200	697	1,059	161
Mean	788	2,311	2,707	396
Maximum	>1,000	2,967	3,285	580

Increase of IOFC (\$/cow per year)

- Between 7 and 52%
- Mean = \$396
- Range = \$161 to \$580

Acknowledgements

This project is supported by Agriculture and Food Research Initiative Competitive Grant No. 2011-68004-30340 from the USDA National Institute of Food and Agriculture



United States Department of Agriculture National Institute of Food and Agriculture

